

## Homework 4 (Stat 620, Fall 2009)

Due Thu Oct 8, in class

- Let  $A(t)$  and  $Y(t)$  denote respectively the age and excess at  $t$ . Find:
  - $\mathbb{P}\{Y(t) > x | A(t) = s\}$ .
  - $\mathbb{P}\{Y(t) > x | A(t + x/2) = s\}$ .
  - $\mathbb{P}\{Y(t) > x | A(t + x) > s\}$  for a Poisson process.
  - $\mathbb{P}\{Y(t) > x, A(t) > y\}$ .
  - If  $\mu < \infty$ , show that, with probability 1,  $A(t)/t \rightarrow 0$  as  $t \rightarrow \infty$ .

Hint: For (d), use a regenerative process argument (E.g. Ross, section 3.7) to find  $\lim_{t \rightarrow \infty} \mathbb{P}(Y(t) > x, A(t) > y)$ . For (e), you may use without proof the following results on convergence with probability 1: **(L1)**  $\lim_{n \rightarrow \infty} S_n/n = \mu$ ; **(L2)**  $\lim_{t \rightarrow \infty} N(t) = \infty$ ; **(L3)**  $\lim_{t \rightarrow \infty} N(t)/t = 1/\mu$ .

- Consider a single-server bank in which potential customers arrive in accordance with a renewal process having interarrival distribution  $F$ . However, an arrival only enters the bank if the server is free when he or she arrives. Would the number of events by time  $t$  constitute a (possibly delayed) renewal process if an event corresponds to a customer:
  - entering the bank?
  - leaving the bank?What if  $F$  were exponential?

- On each bet a gambler, independently of the past, either wins or loses 1 unit with respective probability  $p$  and  $1 - p$ . Suppose the gambler's strategy is to quit playing the first time she wins  $k$  consecutive bets. At the moment she quits
  - find her expected winnings.
  - find the expected number of bets that she has won.

Hint: It may help you to look at Example 3.5(A) in Ross.

- Prove Blackwell's theorem for renewal reward processes. That is, assuming that the cycle distribution is not lattice, show that, as  $t \rightarrow \infty$ ,

$$\mathbb{E}[\text{reward in}(t, t + a)] \rightarrow a \frac{\mathbb{E}[\text{reward incycle}]}{\mathbb{E}[\text{time of cycle}]}$$

Assume that any relevant function is directly Riemann integrable.

Hint: You may adopt an informal approach by assuming that one can write

$$\mathbb{E} \left[ \int_t^{t+a} dR(s) \right] = \int_t^{t+a} \mathbb{E}[dR(s)],$$

and then developing the identity

$$\mathbb{E}[dR(t)] = \mathbb{E}[R_1 | X_1 = t] dF(t) + \int_0^t \{ \mathbb{E}[R_1 | X_1 = t - x] dF(t - x) \} dm(x).$$

If you can find a more elegant or more rigorous solution, that would also be good!

5. The life of a car is a random variable with distribution  $F$ . An individual has a policy of trading in his car either when it fails or reaches the age of  $A$ . Let  $R(A)$  denote the resale value of an  $A$ -year-old car. There is no resale value of a failed car. Let  $C_1$  denote the cost of a new car and suppose that an additional cost  $C_2$  is incurred whenever the car fails.
- (a) Say that a cycle begins each time a new car is purchased. Compute the long-run average cost per unit time.
- (b) Say that a cycle begins each time a car in use fails. Compute the long-run average cost per unit time.
- Note:* In both (a) and (b) you are expected to compute the ratio of the expected cost incurred in a cycle to the expected time of a cycle. The answer should, of course, be the same in both parts.